

Na⁺/H⁺ exchange and its inhibition in cardiac ischemia and reperfusion

W. Scholz and U. Albus

Hoechst AG, Frankfurt/M., FRG

Summary: The characterization of various ion transport systems has led to a better understanding of the effects, which seem to take part in the impairment of ischemic and reperfused cardiac tissue. This review discusses the role of the Na⁺/H⁺ exchange system in the pathophysiology of ischemia and reperfusion and the beneficial effects of its inhibition.

At the onset of ischemia intracellular pH (pH_i) decreases due to anaerobic metabolism and ATP hydrolysis, leading to an activation of Na⁺/H⁺ exchange. This in turn increases intracellular Na⁺ (Na⁺_i) and activates Na⁺/K⁺ ATPase, with a consecutive increase of energy consumption. Since cellular Na⁺ and Ca⁺⁺ transport are coupled by the Na⁺/Ca⁺⁺ exchange system, which depends on the Na⁺ gradient, the high Na⁺_i leads to increased intracellular Ca⁺⁺ (Ca⁺⁺_i). After a certain period, Na⁺/H⁺ exchange is inactivated by a decrease of extracellular pH.

In case of reperfusion the acid extracellular fluid is washed out, which reactivates Na⁺/H⁺ exchange, leading to an unfavourably fast restoration of pH_i and a second time to Na⁺ and Ca⁺⁺_i overflow.

High Ca⁺⁺_i is assumed to be one of the main reasons for ischemic and reperfusion injury, like arrhythmias, myocardial contracture, stunning and necrosis.

It seems that the inhibition of Na⁺/H⁺ exchange can interrupt this process at an early phase and prevent or delay the consequences of ischemia and reperfusion as demonstrated by numerous investigators.

Key words: Na⁺/H⁺ exchange inhibition - cardiac ischemia - reperfusion - intracellular pH - intracellular Na⁺ - intracellular Ca⁺⁺

Introduction

In the last decade it has become clear, that most pathophysiological processes in cardiac ischemia and reperfusion are connected to a derangement of cellular ion homeostasis. The characterization of various ion transport systems has led to the identification of three effects, which seem to play key roles in the impairment of ischemic and reperfused tissue. These interdependent effects are decreased intracellular pH (pH_i), intracellular Na⁺ (Na⁺_i) overload and intracellular Ca⁺⁺ (Ca⁺⁺_i) overload.

The interdependence between intracellular acidosis and elevated Na⁺_i and Ca⁺⁺_i concentrations is based on the activity of certain ion transport systems. Increased activity of the Na⁺ coupled HCO₃⁻ transport and an excessive activation of Na⁺/H⁺ exchange by a decrease of pH_i lead to a significant elevation of Na⁺ influx into the endangered tissue (50, 70, 76, 97). As long as sufficient ATP is available, the intruding Na⁺ ions can be transported against the Na⁺ gradient into the extracellular space by the Na⁺/K⁺ ATPase (25, 75). But eventually, decreasing energy stores and elevated Na⁺ influx result in a marked increase of the concentration of Na⁺_i (25, 46). Since intracellular Na⁺ and Ca⁺⁺ concentrations are linked by a 3Na⁺/Ca⁺⁺ exchange (7, 41, 55), the elevation of Na⁺_i finally causes intracellular Ca⁺⁺ overload (26, 30, 58, 97), which induces cardiac arrhythmias (14, 90) and necrosis (90).

Already in 1984 it has been suggested, that Na^+/H^+ exchange could play an important role in this chain of pathophysiological events (25). While the regulation of pH_i might be dominated by HCO_3^- dependent systems under normal conditions (99), it has been claimed that the exchanger causes a major part of the Na^+ influx in cardiac ischemia and reperfusion (25, 46, 58, 94, 97). More recently, this point of view has been supported by studies showing a significant reduction of Na^+ and Ca^{++} overload (2, 39, 58, 111) and impressive cardioprotective effects with inhibitors of the Na^+/H^+ exchanger (16, 33, 38, 79, 84, 85). Altogether this points to a crucial involvement of this ion transport system in the mechanisms of ischemic and reperfusion injuries.

In the following, this review tries to summarize and interpret the current data on the role of Na^+/H^+ exchange in the pathophysiology of cardiac ischemia and reperfusion. It will then discuss the effects of an inhibition of the exchanger under these conditions.

Role of Na^+/H^+ exchange in ischemia

It seems that there are at least three phases of cardiac ischemia. The first phase would be the time until the ischemic myocardium ceases to contract, the second phase would be the period of ischemic arrhythmias and the third phase would be characterized by a lack of mechanical and electrical activity.

At the beginning of an ischemic insult myocardial cells are subjected to numerous changes in metabolism and ion transport activity. Since the remaining oxygen in the tissue is diminished within few seconds (73) the physiological way of energy production by degrading fatty acids in the mitochondria (104) is inhibited and the production of energy depends on anaerobic glycolysis (1, 92).

However, energy consumption will continue, as for example for the maintenance of ion gradients by Na^+/K^+ ATPase (71). This degradation of ATP is accompanied by a production of protons (17) most of which would be neutralized by the metabolic activity inside the mitochondria under physiological conditions (17). Since mitochondrial activity ceases after few seconds of ischemia (81, 107) the accumulation of protons leads to a decrease of pH_i (59, 61).

The ischemic cells counteract this process by an activation of several pH regulating cellular ion transport systems like lactate transport (100), Na^+ dependent HCO_3^- transport (99) and Na^+/H^+ exchange (25, 73).

There is a controversy about the dominance of the different pH_i regulation systems in ischemia. However, several investigators found that recovery from low pH_i depends on Na^+/H^+ exchange (99, 106) and that the exchanger seems to be the dominant pH-regulating system in acidified cardiac cells (25, 62). Furthermore, it has been shown in acidified cells (25) as well as in isolated hearts that Na^+/H^+ exchange accounts for the majority of Na^+ influx during pH recovery in ischemia and reperfusion (58, 94, 97). Since the activity of the Na^+/K^+ pump is controlled by the concentration of Na^+_i (28), the increased Na^+ influx is followed by an activation of the ATP consuming ion pump. However, degradation of ATP will generate more protons (17) and it will also enhance anaerobic glycolysis (36) and lactate formation. This process creates a vicious cycle that will lead to a further activation of the Na^+ dependent pH regulating systems and to a further increase in Na^+_i .

Even in these first seconds of ischemia degradation of ATP, anaerobic glycolysis, decrease of pH_i , and Na^+ influx are very likely interconnected by several feed-back loops. Decrease of pH_i is caused by anaerobic metabolism and degradation of ATP (17). This drop of pH_i itself causes enhanced Na^+ influx, increased ATP consumption (99, 106), more anaerobic metabolism and consequently a further drop of pH_i . On the other hand, ATP degradation and anaerobic glycolysis as well as numerous intracellular processes depend on pH_i and are

reduced by intracellular acidosis (77). Thus, the cycle of ATP consumption and proton generation can run unhampered only as long as protons can be extruded efficiently to the extracellular space and as long as substantial amounts of ATP are available.

A fast onset of the drop of pH_i and a decrease of ATP of about 50 % within 10 minutes has been observed in ischemic rat hearts at 37°C (23, 27, 58). Within the same time pH_i reached a value of about 6.3.

What are the possible reasons for the observed decrease in pH_i and ATP? Protons, which are generated by the anaerobic metabolism, are transported into the extracellular space. Since they are not washed out during ischemia, the extracellular pH can be expected to decrease. In fact, this has been demonstrated by several investigators (12, 13, 83), and it has been shown that extracellular pH (pH_o) decreases to about 6.0 in ischemic myocardial tissue within 15–20 minutes (23, 24, 27, 42, 57). In consequence, this accumulation of protons in the extracellular space will increasingly inhibit the extrusion of more protons from the ischemic cells (101, 102). In addition, Na^+ -coupled HCO_3^- transport will function only as long as HCO_3^- is available in the extracellular environment, and lactic acid transport will be hampered by a decreasing gradient for lactate and protons across the cell membrane. Consequently, a decrease of pH_i seems inevitable relatively soon after the onset of ischemia.

ATP decreases because anaerobic glycolysis is a much less efficient way of energy production than normal cellular metabolic pathways, yielding only five per cent of the regular amount. In addition, anaerobic glycolysis depends on intracellular glycogen stores, which are depleted during ischemia, because of lacking glucose supply in the ischemic tissue. Another reason is the fall of pH_i itself, which inhibits anaerobic glycolysis like most other enzyme-driven pH dependent cellular reactions (60, 77, 78).

There are several consequences of decreased pH_i and ATP. One consequence is a decrease of myocardial contractility, which is observed very soon after the beginning of ischemia (11, 31, 52, 53, 63, 66, 91, 92, 103). Since ATP is required for the contractile process, a reduction of myocardial contraction can be expected with ATP deficiency. However, the reported drop of ATP of only about 10–15 % (10, 29, 109) in this first phase of ischemia does not well correlate with the fast reduction of myocardial contraction. In contrast, the faster decrease of pH_i (13, 20, 57) gives a much better correlation. This is in agreement with the diminished binding of Ca^{++} to the contractile elements in acidified myocardial cells (8, 22, 53) as well as the reduced activity of the myosin-ATPase at low pH (8, 40, 80). In addition, myocardial contractility seems to be very sensitive to only small reductions of pH_i (57).

As reported earlier, pH_i regulating ion transport systems cause Na^+ influx (25, 46, 71, 100). However, a decrease of intracellular ATP will reduce the capacity of the ATP dependent Na^+/K^+ ATPase to eliminate Na^+ from the intracellular space. In fact, a marked increase of Na^+_i in ischemic cardiac tissue has been observed by numerous investigators (48, 50, 58, 70, 76, 97). In parallel with the increase of Na^+ in ischemic cardiac tissue an increase of intracellular Ca^{++} has been found (58,94). The Ca^{++} levels in myocardial cells are significantly regulated by $3\text{Na}^+/\text{Ca}^{++}$ exchange (7, 41, 55). This exchanger is driven by the electrical potential of the cell and by the Na^+ and Ca^{++} gradient (7, 41, 55) and it extrudes Ca^{++} under physiological conditions (7, 41, 55). The transport rate of the $3\text{Na}^+/\text{Ca}^{++}$ exchange is diminished by intracellular Na^+ overload (26, 30, 46, 47, 69, 74, 97) and the exchanger might even change the direction of its transport if Na^+_i becomes excessively high (25, 46, 98).

High Ca^{++}_i levels are thought to be deleterious to ischemic myocardium in several ways (14, 26, 90). Together with low ATP high Ca^{++}_i has been connected with overexcitability and electrical instability (4). In general, it is well documented that high Ca^{++}_i is one of the main reasons for ischemic arrhythmias (and reperfusion arrhythmias) (65).

Cardiac arrhythmias usually do not start immediately at the beginning of an ischemic insult. In patients undergoing Percutaneous Transluminal Coronary Angioplasty (PTCA) very few arrhythmias are observed during the one or two minutes of cardiac ischemia, which the patients have to suffer (64). In anesthetized rats with experimental ligation of the left coronary artery marked arrhythmic activity starts after 5 minutes of ischemia and continues for about 10 more minutes (84, 85). These events might reflect a certain degree of cellular Na^+ and Ca^{++} overload after 5 minutes of ischemia. Indeed, NMR measurements in rat hearts show significant increases of Na^+_i (35, 70) and Ca^{++}_i (23) after this period of ischemia.

It may also be interesting to consider the pathophysiological process in adrenergic nerve endings in the heart during ischemia and reperfusion. One of the significant activities of these cells is the uptake of norepinephrine from the extracellular space by a Na^+ driven uptake mechanism (uptake₁) (87, 96). Inside the cells norepinephrine is transported into storage vesicles in exchange for protons (6, 68, 108).

During an ischemic insult adrenergic nerve endings very likely undergo a similar derangement of metabolism and ion homeostasis as myocardial cells (88). There should also be a decrease of pH_i and an increase of Na^+ influx through Na^+/H^+ exchange, leading to an intracellular Na^+ overload and a decrease of ATP (3, 87, 88). It has been shown that the activity of the Na^+ -dependent norepinephrine uptake₁ is diminished by high Na^+_i and that it might also change its direction of transport (87).

Furthermore, the storage mechanism of norepinephrine in the vesicles depends on energy. It will become insufficient when ATP decreases in ischemia (6, 68, 108), leading to intracellular norepinephrine overflow (6, 68, 108). A high intracellular norepinephrine concentration together with high Na^+ will cause an extracellular overflow of norepinephrine through reversed transport of the uptake₁ (87, 96). Under such conditions the extracellular norepinephrine concentration in cardiac tissue might rise a hundred fold above normal (87). Since the arrhythmogenic potential of high norepinephrine concentrations is well known (15, 49), norepinephrine overflow might be one of the arrhythmogenic factors in ischemic cardiac tissue (88).

Arrhythmias in cardiac ischemia cease after 10 to 15 min in the rat model, although Na^+ and Ca^{++} overload continue to increase beyond that time (58, 94). Furthermore, pH_i decreases down to about 6.0 after 20–30 min (23, 24, 27, 35, 42, 57), which is accompanied by a pronounced decrease of ATP in the same time (23, 27, 35, 94). In this respect, it is remarkable that neither pH_i nor pH_o tend to drop below a value of about 6.0, even if ischemia lasts longer than 15–30 min (23, 35).

This lack of further acidification could indicate a stop of a significant metabolism in the ischemic myocardial cells. Anaerobic glycolysis, which is known to depend on pH_i (60, 77, 78), is very likely inhibited when pH_i reaches values as low as 6.0. Without energy the maintenance of ion gradients, necessary for the electrical potentials of the cells, is impossible. Therefore, ischemic cells become electrically quiescent (43) and arrhythmic activity will stop in the ischemic core area.

Although intracellular Ca^{++} overload (58, 94) cannot induce further arrhythmic activity it will, however, destroy mitochondria (9, 89, 90), activate lysosomes (45), destabilize the cell membrane (32, 90), and finally cause cellular necrosis (90). Without reperfusion this might be the inevitable fate of the endangered cells.

Role of Na^+/H^+ exchange in reperfusion

Reperfusion, for the rescue of the ischemic tissue, is a double-edged sword. In the beginning it causes a very rapid washout of the acidic extracellular fluid in the ischemic area.

This generates a huge pH-gradient between intracellular and extracellular space. In consequence, the pH-regulating cellular ion transport systems are reactivated to a maximum degree, which will cause a large Na^+ influx by Na^+/H^+ exchange (46) and Na^+ dependent HCO_3^- transport (100).

The Na^+ ions cannot be extruded sufficiently by the Na^+/K^+ ATPase due to a lack of ATP, leading to an additional intracellular Na^+ overload (97) and, secondary, to an additional Ca^{++} overload (93) by $3\text{Na}^+/\text{Ca}^{++}$ exchange (58, 95). This process is paralleled by a very fast restoration of pH_i (12, 23, 100). While elevated Ca^{++} in myocardial cells is less efficacious at low pH_i (8, 22, 53) it will increasingly take effect when normal pH_i is restored. The results are reperfusion arrhythmias (65), reperfusion injuries like myocardial contraction (5, 94), myocardial stunning (5, 42, 94), and necrosis (5, 72, 94).

Energy depleted local sympathetic nerve endings face the same process as ischemic myocardium during reperfusion (87). In the nerve endings Na^+ overload and ATP depletion again result in norepinephrine overflow by the above described mechanism (87). This perpetuates arrhythmias (15, 67) and enhances Ca^{++} influx (104) in the destabilized myocardial tissue.

Adverse effects of reperfusion seem to be limited to an early phase. Reperfusion experiments with acidic solutions, which keep Na^+/H^+ exchange and Na^+ coupled HCO_3^- transport inactivated, have shown, that this vulnerable phase in cardiac reperfusion might last for about two minutes (4). While two minutes of acidic reperfusion had a maximum protective effect in cardiac tissue, reperfusion, continued with physiological pH after two minutes, had no further adverse effects (4). This finding suggests, that after two minutes of reperfusion without further Na^+ overflow and a delayed restoration of pH_i , intracellular ATP is restored so far that the reperfused cells might cope with enhanced Na^+ influx and might reduce elevated Ca^{++} .

In summary, Na^+/H^+ exchange seems to play a key role in the pathophysiology of ischemia and reperfusion in multiple ways, and beneficial effects from Na^+/H^+ exchange inhibition in cardiac ischemia and reperfusion can be expected.

Inhibition of Na^+/H^+ exchange in ischemia

In ischemia cardiac cells rapidly shift to anaerobic metabolism followed by activation of Na^+/H^+ exchange (73). Therefore, one of the first significant effects of Na^+/H^+ exchange inhibition in ischemia should be a reduction of cellular Na^+ influx (46). This might have several consequences (Figs. 1a and 1b). Primarily, there would be less Na^+ ions to be extruded by the Na^+/K^+ pump. Secondly, the ATP consumption by the Na^+/K^+ pump would be reduced and Na^+ overload would be prevented or at least delayed. The next consequence would be a reduction of the following intracellular Ca^{++} overload via the $3\text{Na}^+/\text{Ca}^{++}$ exchange (54). Furthermore, a prevention of Ca^{++} overload should prevent ischemic arrhythmias if ischemic arrhythmias result from increased Ca^{++} .

Most of these conclusions are already supported by experimental evidence. In fact, a marked reduction of Na^+ overload and a conservation of ATP (84, 85) by Na^+/H^+ exchange inhibition in cardiac ischemia has been observed by several investigators (2, 51, 111). In addition, it seems that Na^+/H^+ exchange inhibition causes a diminution of Ca^{++} overload (2, 39, 58, 111). Furthermore, ischemic arrhythmias can almost completely be prevented by Na^+/H^+ exchange inhibitors (79, 84, 85).

Since the exchanger has a significant role in the regulation of pH_i in cardiac ischemia (99, 100), the decrease of pH in ischemic myocardial cells should be accelerated by Na^+/H^+ exchange inhibition. This would enhance the inhibition of anaerobic glycolysis (60, 77, 78) and reduce the degradation of ATP (77). In addition, less protons would be generated by




Fig. 1a. Active Na^+/H^+ exchange in ischemic cardiac myocytes: Elevated intracellular Na^+ concentrations cause an intracellular Ca^{++} overload. Anaerobic metabolism is driven by high intracellular Na^+ and the excretion of H^+ .




Fig. 1b. Inhibited Na^+/H^+ exchange in ischemic cardiac myocytes: Reduced intracellular Na^+ concentrations prevent intracellular Ca^{++} overload. Anaerobic metabolism is inhibited by reduced intracellular Na^+ and intracellular acidification.

ATP break down, since a decreased Na^+ influx would reduce ATP consumption by Na^+/K^+ ATPase.

At present, there is only few experimental evidence for an influence of an inhibition of the exchanger on the pH_i decrease in ischemia. While the decrease of pH_i seemed slightly faster

in ischemic ferret hearts pre-treated with the Na^+/H^+ exchange inhibitor EIPA (100), another investigator did not find any effect of amiloride on the rate of fall of pH_i in rat hearts (58). However, the shortcoming of most NMR studies using the phosphate shift to investigate changes of pH_i in ischemia and reperfusion is the relatively infrequent data acquisition of pH like every 5 or 10 min. Thus, an early difference in pH_i by Na^+/H^+ exchange inhibition would not necessarily be recognized.

On the other hand, Na^+/H^+ exchange inhibition causes a diminution of lactate efflux from ischemic cardiac tissue as well as a decrease of cellular lactate content (84, 85). These observations suggest a decrease of lactate formation, which would implicate a reduction of anaerobic glycolysis. In the same studies these effects were paralleled by a conservation of intracellular glycogen stores (84, 85). Since glucose has to come from intracellular glycogen stores during ischemia, this observation also indicates a reduction of anaerobic glycolysis by Na^+/H^+ exchange inhibition.

A faster decrease of pH_i by Na^+/H^+ exchange inhibition in ischemia could have further beneficial effects. One of them could be a reduced activity of the $3\text{Na}^+/\text{Ca}^{++}$ exchange which has been found to depend on pH_i (21, 37). Another beneficial effect might be a reduction of free radical formation, which is also depressed by low pH_i (110). A third one might be an earlier onset of the decrease of contractility of ischemic myocardial tissue. This has been shown with EIPA (83) and might result in a conservation of cellular energy stores. However, there are only few experimental data available yet on these possible effects of Na^+/H^+ exchange inhibition.

A conservation of cellular energy stores, as observed with Na^+/H^+ exchange inhibition in myocardial tissue might also prevent or diminish norepinephrine overflow in ischemia from intracellular vesicles in the local sympathetic nerve endings (88). Furthermore, a reduced Na^+_i overload (88) might avert the release of norepinephrine by a reversed uptake₁ mechanism (88).

In fact, such a prevention of norepinephrine overflow in ischemic cardiac tissue has been demonstrated in several experiments with different Na^+/H^+ exchange inhibitors (87). These effects in neuronal tissue in the heart might contribute to the reduction of Ca^{++}_i overload and arrhythmic activity in cardiac ischemia, which has been observed with Na^+/H^+ exchange inhibitors.

As mentioned above, in prolonged ischemia, when arrhythmic activity has finally ceased, a continuing Ca^{++}_i overload will cause irreversible injury and finally necrosis of the ischemic cardiac tissue (90) by damaging mitochondria (9, 89, 90), activating lysosomes (45) and destabilizing cellular membranes (32, 90).

Na^+/H^+ exchange inhibitors might prevent or delay this process by a marked reduction of Ca^{++}_i in ischemic cardiac tissue (58). This assumption has been partially confirmed by studies using isolated ischemic working rat hearts. In these experiments Na^+/H^+ exchange inhibitors markedly decreased the release of the intracellular enzymes creatine kinase and lactate dehydrogenase, indicating a reduction of cellular membrane instability and necrosis (84, 85).

It is hard to predict, how long the protection by Na^+/H^+ exchange inhibition could last. In human hearts "hibernating myocardium" has been identified with positron emission tomography after relatively long periods of reduced flow (82). Although there is not much known about the processes involved, this observation demonstrates that a protection of underperfused cardiac tissue for days or even weeks might be possible. Nevertheless, the best chance of the ischemic myocardium for survival and restoration of contractile function is reperfusion. The benefits of reperfusion might be markedly increased, if reperfusion injury could be diminished.

Inhibition of Na^+/H^+ exchange in reperfusion

As described above, the sudden removal of the acidic extracellular fluid during reperfusion leads to a second excessive Na^+ influx (4). Since both Na^+ influx and the fast rise in pH_i are mostly caused by Na^+/H^+ exchange, an inhibition of the exchanger in the vulnerable phase of reperfusion might significantly reduce reperfusion injuries.

A decrease of Na^+ influx during reperfusion would again reduce ATP consumption by the Na^+/K^+ ATPase (46) and would prevent Na^+_i and Ca^{++}_i overflow (25, 46). In parallel, a retarded recovery of pH_i would protect the reperfused cardiac tissue from the effects of the high Ca^{++}_i concentrations (8, 22, 40, 53, 80) gained during ischemia (93).

Ischemia-reperfusion experiments with Na^+/H^+ exchange inhibitors have already demonstrated some beneficial effects like a marked reduction of intracellular Na^+ and Ca^{++} overload in reperfused myocardium (83), a slow down of pH_i recovery (100), increased ATP levels (84, 85), and a prevention of norepinephrine overflow (87).

In consequence, all these actions of Na^+/H^+ exchange inhibition in reperfusion seem to have a common outcome, which is the observed prevention of intracellular Ca^{++} overload (58). Since excessively elevated intracellular Ca^{++} is thought to be the reason for adverse effects of reperfusion, like arrhythmias (14, 26, 90), myocardial contracture (5, 94), myocardial stunning (5, 94), and necrosis (5, 72, 94), a prevention of these effects by Na^+/H^+ exchange inhibition could be expected. In fact, a diminution of reperfusion arrhythmias with Na^+/H^+ exchange inhibitors has been demonstrated recently in vitro and in vivo by numerous investigators (16, 38, 79, 83–85). In parallel, a prevention of the increase of left ventricular enddiastolic pressure (18, 33, 58), a prevention of myocardial stunning in isolated hearts (33) and in anesthetized pigs (79), and a reduction of myocardial damage (38) have been observed.

Besides a reduction of intracellular Na^+ and Ca^{++} overload in reperfused cardiac tissue (58), Na^+/H^+ exchange inhibitors could be even more protective if they are also present during the ischemic period. Cardiac tissue might be in a better condition then to meet reperfusion stress. At the time being, only few studies have been reported in which Na^+/H^+ exchange inhibitors were employed during both, ischemia and reperfusion, or during reperfusion only (38). However, recent experiments with blood perfused rabbit hearts, using the highly specific Na^+/H^+ exchange inhibitor HOE 694 (84), clearly demonstrated a better prevention of myocardial contracture and a better recovery of contractility when the inhibitor was present during ischemia and reperfusion (33).

Inhibition of Na^+/H^+ exchange in low flow ischemia, in hypoxia and in reperfusion after low flow ischemia

Experiments under so-called "low flow ischemia" gave controversial results about the involvement of the Na^+/H^+ exchanger. This variance probably derives from different degrees of ischemia reached in the different experiments.

While in one study, with isolated nonworking rat hearts, under low flow conditions only a small decrease of pH_i and ATP occurred, and the Na^+/H^+ exchange inhibitor EIPA had no effect on intracellular ATP or pH (34), another study showed a marked decrease of intracellular pH under hypoxic conditions (105). Other investigators have found a decrease of intracellular Na^+ and Ca^{++} overload by Na^+/H^+ exchange inhibition (2).

Two more studies reported significant protection by amiloride derivatives against reperfusion injury after low flow ischemia (16, 18).

Conclusion

While the part of Na^+/H^+ exchange in the regulation of pH_i might be negligible under normal conditions (99), the exchanger is markedly activated in ischemia and reperfusion (25,

46, 71, 100). Furthermore, evidence has been accumulated since 1985 for a multiple involvement of Na^+/H^+ exchange in the pathophysiology of cardiac ischemia and reperfusion.

Several studies have demonstrated that the inhibition of the exchanger in cardiac ischemia and reperfusion definitely decreases Na^+_i and Ca^{++}_i overload (2, 39, 58, 83, 111). In addition, inhibitors of Na^+/H^+ exchange cause a reduction of anaerobic glycolysis (84, 85) and diminish ATP depletion (51, 84–86).

As expected, these effects were paralleled by a reduction of arrhythmias and necrosis in cardiac ischemia (19, 56), as well as by a prevention of reperfusion injury (18, 33, 38, 58, 79, 85). These observations fit elegantly to the assumed significant role of the exchanger in the pathophysiology of ischemia and reperfusion. However, their conclusiveness depends on the specificity of the actions of the Na^+/H^+ exchange inhibitors used on the exchange system.

While amiloride, as the first substance found to affect Na^+/H^+ exchange, has qualified as a relatively weak and non-specific inhibitor (44), several potent and rather selective amiloride derivatives, like EIPA, DMA, MIBA, HMA and others have been identified in the last decade (44). In addition, a new and highly specific compound, HOE 694, which has been characterized as rather potent and selective inhibitor of the exchanger, has been employed in several studies (33, 79, 84). The protective effects, which all these compounds have shown in cardiac ischemia and reperfusion, are comparable in their principle quality. Therefore, these effects seem to be definitely related to Na^+/H^+ exchange inhibition.

In summary, pathophysiological evidence and experimental observations have qualified Na^+/H^+ exchange inhibition as a possible new therapeutic principle in cardiac ischemia and reperfusion.

References

1. Allen DG, Orchard CH (1987) Myocardial contractile function during ischemia and hypoxia. *Circulat Res* 60:153–168
2. Anderson SE, Murphy E, Steenbergen C, London RE, Cala PM (1990) Na-H exchange in myocardium: effects of hypoxia and acidification on Na and Ca. *Am J Physiol* 259:C940–C948
3. Aronson PS (1985) Kinetic properties of the plasma membrane Na^+/H^+ -exchanger. *Ann Rev Physiol* 47:545–560
4. Avkiran M, Ibuki C (1992) Reperfusion-induced arrhythmias – a role for washout of extracellular protons? *Circulat Res* 71:1429–1440
5. Barry WH, Peeters GA, Rasmussen CAF, Cunningham MJ (1987) Role of changes in $[\text{Ca}^{++}]_i$ in energy deprivation contracture. *Circulat Res* 61:726–734
6. Beers MF, Carty SE, Johnson RG, Scarpa A (1982) H^+ -ATPase and catecholamine transport in chromaffin granules. *Ann NY Acad Sci* 402:116–133
7. Beuckelmann DJ, Wier WG (1989) Sodium-calcium exchange in guinea-pig cardiac cells: Exchange current and changes in intracellular Ca^{++} . *J Physiol* 414:499–520
8. Blanchard EM, Solaro RJ (1984) Inhibition of the activation and troponin calcium binding of dog cardiac myofibrils by acidic pH. *Circulat Res* 55:382–391
9. Borgers M, Piper HM (1986) Calcium shifts in anoxic cardiac myocytes: A cytochemical study. *J Mol Cell Cardiol* 18:439–448
10. Camacho SA, Lanzer P, Toy J, Gober J, Valenza M, Botvinick EH, Weiner MW (1988) In vivo alterations of high-energy phosphates and intracellular pH during reversible ischemia in pigs: A ^{31}P magnetic resonance spectroscopy study. *Amer Heart J* 116:701–708
11. Cingolani HE, Mattiazzi AR, Blesa ES, Gonzalez HE (1970) Contractility in isolated mammalian heart muscle after acid-base changes. *Circulat Res* 26:269–278
12. Cobbe SM, Poole-Wilson PA (1980) Tissue acidosis in myocardial hypoxia. *J Mol Cell Cardiol* 12:761–770
13. Cobbe SM, Poole-Wilson PA (1980) The time of onset and severity of acidosis in myocardial ischaemia. *J Mol Cell Cardiol* 12:745–760
14. Coetzee WA, Opie LH (1987) Effects of components of ischemia and metabolic inhibition on delayed afterdepolarizations in guinea pig papillary muscle. *Circulat Res* 61:157–165

15. Corr PB, Gillis RA (1978) Autonomic neutral influences on the dysrhythmias resulting from myocardial infarction. *Circulat Res* 43:1-9
16. Dennis SC, Coetzee WA, Cragoe EJ, Opie LH (1990) Effects of proton buffering and of amiloride derivatives on reperfusion arrhythmias in isolated rat hearts. *Circulat Res* 66:1156-1159
17. Dennis SC, Gevers W, Opie LH (1991) Protons in ischemia: Where do they come from; Where do they go to? *J Mol Cell Cardiol* 23:1077-1086
18. Duan J, Karmazyn M (1992) Protective effects of amiloride on the ischemic reperfused rat heart. Relation to mitochondrial function. *Europ J Pharmacol* 210:149-157
19. Duff HJ, Lester WM, Rahmberg M (1988) Amiloride, antiarrhythmic and electrophysiological activity in the dog. *Circulation* 78:1469-1477
20. Elliot AC, Smith GL, Eisner DA, Allen DG (1992) Metabolic changes during ischaemia and their role in contractile failure in isolated ferret hearts. *J Physiol* 454:467-490
21. Ellis D, MacLeod KT (1985) Sodium-dependent control of intracellular pH in Purkinje fibers of sheep heart. *J Physiol* 359:81-105
22. Fabiato A, Fabiato F (1978) Effects of pH on the myofilaments and the sarcoplasmic reticulum of skinned cells from cardiac and skeletal muscle. *J Physiol* 276:233-255
23. Fralix TA, Murphy E, London RE, Steenbergen C (1993) Protective effects of adenosine in the perfused rat heart: changes in metabolism and intracellular ion homeostasis. *Am J Physiol* 264:C986-C994
24. Fralix TA, Steenbergen C, London RE, Murphy E (1993) Glibenclamide does not abolish the protective effect of preconditioning on stunning in the isolated perfused rat heart. *Cardiovasc Res* 27:630-637
25. Frelin C, Vigne P, Lazdunski M (1984) The role of Na^+/H^+ exchange system in cardiac cells in relation to the control of the internal Na^+ concentration. *J Biol Chem* 259:8880-8885
26. Frelin C, Vigne P, Lazdunski M (1985) The role of Na^+/H^+ exchange system in the regulation of the internal pH in cultured cardiac cells. *Eur J Biochem* 149:1-4
27. Garlick PB, Radda GK, Seeley PJ (1979) Studies of acidosis in the ischemic heart by phosphorus nuclear magnetic resonance. *Biochem J* 184:547-554
28. Glitsch HG (1982) Electrogenic Na pumping in the heart. *Ann Rev Physiol* 44:389-400
29. Gudbjarnason S, Mathes P, Ravens KG (1970) Functional compartmentation of ATP and creatine phosphate in heart muscle. *J Mol Cell Cardiol* 1:325-339
30. Haigney MCP, Miyata H, Lakatta EG, Stern MD, Silverman HS (1992) Dependence of hypoxic cellular calcium loading on $\text{Na}^+/\text{Ca}^{++}$ exchange. *Circulat Res* 71:547-557
31. Harrison SM, Frampton JE, McCall E, Boyett MR, Orchard CH (1992) Contraction and intracellular Ca^{++} , Na^+ , and H^+ during acidosis in rat ventricular myocytes. *Am J Physiol* 262:C348-C357
32. Hearse DJ, Humphrey SM, Bullock GR (1978) The oxygen paradox and the calcium paradox: two facets of the same problem? *J Mol Cell Cardiol* 10:641-668
33. Hendrikx M, Mubagwa K, Flameng W (1993) Inhibition of Na^+/H^+ exchange protects against postischemic dysfunction in isolated, blood perfused rabbit heart. *J Mol Cell Cardiol* 25:VI P 12 (Abstract)
34. Imai S, Shi AY, Ishibashi T, Nakazawa M (1991) Na^+/H^+ exchange is not operative under low flow ischemic conditions. *J Mol Cell Cardiol* 23:505-517
35. Ishibashi T, Nakazawa M, Imai S (1993) Ischemic changes in myocardial ionic contents of the isolated perfused rat heart as studies by NMR. *Molec Cell Biochem* 119:109-120
36. Jungermann K, Möhler H (1980) *Biochemie*. Springer-Verlag, Berlin Heidelberg New York pp 165-167
37. Kaila K, Vaughan-Jones RD (1987) Influence of sodium-hydrogen exchange on intracellular pH, sodium and tension in sheep cardiac Purkinje fibers. *J Physiol* 390:93-118
38. Karmazyn M (1988) Amiloride enhances postischemic ventricular recovery: possible role of Na^+/H^+ exchange. *Am J Physiol* 255:H608-H615
39. Kazumasa H, Franklin A, Johnson RG, Grossman W, Morgan JP (1992) Na^+/H^+ ; $\text{Na}^+/\text{Ca}^{++}$ exchange enhance and ATP-sensitive K^+ channels ameliorate cell Ca^{++} rise in ischemia. *Circulation* 86:1903 (Abstract)
40. Kentish JC, Nayler WG (1979) The influence of pH on the Ca^{2+} -regulated ATPase of cardiac and white skeletal myofibrils. *J Mol Cell Cardiol* 11:611-617

41. Kimura J, Noma A, Irisawa H (1986) Na-Ca exchange current in mammalian heart cells. *Nature* 319:596-597
42. Kitakaze M, Weisfeldt ML, Marban E (1988) Acidosis during early reperfusion prevents myocardial stunning in perfused ferret hearts. *J Clin Invest* 82:920-927
43. Kleber AG (1983) Resting membrane potential, extracellular potassium activity, and intracellular sodium activity during acute global ischemia in isolated perfused guinea pig hearts. *Circulat Res* 52:442-450
44. Kleyman TR, Cragoe EJ (1988) Antiarrhythmic activity of amiloride: mechanisms. *J Membr Biol* 105:1-21
45. Kutryk MJB, Dhalla NS (1988) Alterations in cardiac lysosomal hydrolases following induction of the calcium paradox. *Can J Physiol Pharmacol* 65/11:2175-2181
46. Lazdunski M, Frelin C, Vigne P (1985) The sodium/hydrogen exchange system in cardiac cells: Its biochemical and pharmacological properties and its role in regulating internal concentrations of sodium and internal pH. *J Mol Cell Cardiol* 17:1029-1042
47. Lederer WJ, Niggli E, Hadley RW (1990) Sodium-calcium exchange in excitable cells: Fuzzy space. *Science* 248:283
48. MacLeod KT (1989) Effects of hypoxia and metabolic inhibition on the intracellular sodium activity of mammalian ventricular muscle. *J Physiol* 416:455-468
49. Malliani A, Schwartz PJ, Zanchetti A (1980) Neural mechanisms in life-threatening arrhythmias. *Amer Heart J* 100:705-715
50. Malloy CR, Buster DC, Margarida M, Castro M, Geraldine CFGC, Jeffrey FMH, Sherry AD (1990) Influence of global ischemia on intracellular sodium in the perfused rat heart. *Magn Reson Med* 15:33-44
51. Maron R (1989) Functional and metabolic protection by Na^+/H^+ exchange inhibition in global ischemia. *J Mol Cell Cardiol* 21:1226 (Abstract)
52. Mattiazzi AR, Cingolani HE (1977) Biphasic effect of hypercapnia on myocardial contractility. *Arch Int Phys Biochem* 85:11-25
53. Mattiazzi AR, Cingolani HE, Spacapan de Castuma E (1979) Relationship between calcium and hydrogen ions in heart muscle. *Am J Physiol* 237:H497-H503
54. Meng HP, Lonsberry BB, Pierce GN (1991) Influence of perfusate pH on the postischemic recovery of cardiac contractile function: Involvement of sodium-hydrogen exchange. *J Pharmacol Exp Ther* 258:772-777
55. Miura Y, Kimura J (1989) Sodium-calcium exchange current. Dependence on internal Ca^{++} and Na^+ and competitive binding of external Na and Ca^{++} . *J Gen Physiol* 93:1129-1145
56. Moffat MP (1989) Amiloride modifies the response of canine Purkinje fibres to conditions of ischemia and reperfusion. *J Mol Cell Cardiol* 21:S143 (Abstract)
57. Mohabir R, Lee HC, Kurz RW, Clusin WT (1991) Effects of ischemia and hypercarbic acidosis on myocyte calcium transients, contraction, and pH_i in perfused rabbit hearts. *Circulat Res* 69:1525-1537
58. Murphy E, Perlman M, London RE, Steenbergen C (1991) Amiloride delays the ischemia induced rise in cytosolic free calcium. *Circulat Res* 68:1250-1258
59. Neely JR, Feuvray D (1981) Metabolic products and myocardial ischemia. *Am J Physiol* 102:282-291
60. Neely JR, Morgan HE (1974) Relationship between carbohydrate and lipid metabolism and the energy balance of heart muscle. *Ann Rev Physiol* 36:413-459
61. Neely JR, Rovetto MJ, Whitmer JT, Morgan HE (1973) Effects of ischemia on function and metabolism of the isolated working rat heart. *Am J Physiol* 225:651-658
62. Ng LL, Davies JE, Quinn P (1993) Intracellular pH regulation in isolated myocytes from adult rat heart in HCO_3^- -containing and HCO_3^- -free media. *Clin Sci* 84:133-139
63. Ng ML, Levy MN, Zieske HA (1967) Effects of changes of pH and carbon dioxide tension on left ventricular performance. *Am J Physiol* 213:115-120
64. Oh JK, Shub C, Ilstrup DM, Reeder GS (1985) Creatine kinase release after successful percutaneous transluminal coronary angioplasty. *Amer Heart J* 109:1225-1231
65. Opie LH, Coetsee WA (1988) Role of calcium ions in reperfusion arrhythmias: Relevance to pharmacologic intervention. *Cardiovasc Drugs Ther* 2:623-636

66. Pannier JL, Leusen I (1968) Contraction characteristics of papillary muscle during changes in acid-base composition of the bathing-fluid. *Arch Int Phys Biochem* 76:624-634
67. Penny WJ (1984) The deleterious effects of myocardial catecholamines on cellular electrophysiology and arrhythmias during ischemia and reperfusion. *Eur Heart J* 5:960-973
68. Phillips JH (1982) Dynamic aspects of chromaffin granule structure. *Neuroscience* 7:1595-1609
69. Pierce GN, Maddaford TG, Kroeger EA, Cragoe EJ (1990) Protection by benzamil against dysfunction and damage in rat myocardium after calcium depletion and repletion. *Am J Physiol* 258:H17-H23
70. Pike MM, Kitakaze M, Marban E (1990) ^{23}Na -NMR measurements of intracellular sodium in intact perfused ferret hearts during ischemia and reperfusion. *Am J Physiol* 259:H1767-H1773
71. Piwnica-Worms D, Jacob R, Shigeto N, Horres CR, Lieberman M (1986) Na/H exchange in cultured chick heart cells: Secondary stimulation of electrogenic transport during recovery from intracellular acidosis. *J Mol Cell Cardiol* 18:1109-1116
72. Poole-Wilson PA (1984) Therapeutic approaches to myocardial infarct size limitation. In: Hearse DJ, Yellon DM (eds) What causes cell death. Raven press, New York, pp 43-60
73. Poole-Wilson PA (1989) Regulation of intracellular pH in the myocardium; relevance to pathology. *Molec Cell Biochem* 89:151-155
74. Rabkin SW (1989) Comparison of the effect of amiloride and its analogue dichlorobenzamil on cardiac chronotropic responses to Ouabain in myocardial cell aggregates in culture. *Pharmacology* 39:230-239
75. Rasmussen HH, Cragoe EJ, Ten Eick RE (1989) Na^+ -dependent activation of Na^+ - K^+ pump in human myocardium during recovery from acidosis. *Am J Physiol* 256:H256-H264
76. Regan TJ, Broisman L, Haider B, Eaddy C, Oldewurtel A (1980) Dissociation of myocardial sodium and potassium alterations in mild versus severe ischemia. *Am J Physiol* 238:H575-H580
77. Rouslin W (1983) Protonic inhibition of the mitochondrial oligomycin-sensitive adenosine 5'-triphosphatase in ischemic and autolyzing cardiac muscle. *J Biol Chem* 258:9657-9661
78. Rouslin W, Erickson JL (1986) Factors affecting loss of mitochondrial functioning in autolyzing cardiac muscle. *J Mol Cell Cardiol* 18:1187-1195
79. Sack S, Mohri M, Schwarz ER, Arras M, Schaper J, Schaper W (1992) Inhibition of Na^+ /H $^+$ exchange prevents ventricular fibrillation and preserves function in porcine stunned myocardium. *J Mol Cell Cardiol* 24:S92 (Abstract)
80. Schadler M (1967) Proportionale Aktivierung von ATPase-Aktivität und Kontraktionsspannung durch Calciumionen in isolierten kontraktile Strukturen verschiedener Muskelarten. *Pfluegers Arch* 296:70-90
81. Schaffer SW, Safer B, Ford C, Illingworth J, Williamson JR (1978) Respiratory acidosis and its reversibility in perfused rat heart: regulation of citric acid cycle activity. *Am J Physiol* 234:H40-H51
82. Schelbert HR, Buxton D (1988) Insights into coronary artery disease gained from metabolic imaging. *Circulat Res* 78:496-505
83. Scheufler E, Henrichs M, Guttman I, Mozes A, Wilffert B (1993) Effect of the Na/H exchange inhibitor Ethyl-Isopropyl-Amiloride (EIPA) during ischaemia and reperfusion. *Brit J Pharmacol* 108:118P
84. Scholz W, Albus U, Lang HJ, Linz W, Martorana PA, Englert HC, Schölkens BA (1993) Hoe 694, a new Na^+ /H $^+$ exchange inhibitor and its effects in cardiac ischaemia. *Brit J Pharmacol* 109:562-568
85. Scholz W, Albus U, Linz W, Martorana P, Lang HJ, Schölkens BA (1992) Effects of Na^+ /H $^+$ exchange inhibitors in cardiac ischemia. *J Mol Cell Cardiol* 24:731-739
86. Scholz W, Albus U, Linz W, Schölkens BA (1991) Effect of Na^+ /H $^+$ exchange inhibition in cardiac ischemia. *Eur Heart J* 12 (Suppl):124 (Abstr)
87. Schömig A, Kurz T, Richardt G, Schömig E (1988) Neuronal sodium homeostasis and axoplasmic amine concentration determine calcium-independent noradrenaline release in normoxic and ischemic rat heart. *Circulat Res* 63:214-226
88. Schömig A, Richardt G (1990) The role of catecholamines in ischemia. *J Cardiovasc Pharmacol* 16:S105-S112
89. Sharma GP, Varley KG, Kim SW, Barwinsky J, Cohen M, Dhalla NS (1975) Alterations in energy metabolism and ultrastructure upon reperfusion of the ischemic myocardium after coronary occlusion. *Am J Cardiol* 36:234-243

90. Shen AC, Jennings RB (1972) Myocardial calcium and magnesium in acute ischemic injury. *Am J Pathol* 67:417-440
91. Smith HW (1926) The actions of acids on turtle muscle with reference to the penetration of anions. *Am J Physiol* 76:411-447
92. Steenbergen C, Deleew G, Rich T, Williamson JR (1977) Effects of acidosis and ischemia on contractility and intracellular pH of rat heart. *Circulat Res* 41:849-858
93. Steenbergen C, Murphy E, Levy L, London RE (1987) Elevation in cytosolic free calcium concentration early in myocardial ischemia in perfused rat heart. *Circulat Res* 60:700-707
94. Steenbergen C, Murphy E, Watts JA, London RE (1990) Correlation between cytosolic free calcium, contracture, ATP, and irreversible ischemic injury in perfused rat heart. *Circulat Res* 66:135-146
95. Steenbergen C, Perlman ME, London RE, Murphy E (1993) Mechanisms of preconditioning: ionic alterations. *Circulat Res* 72:112-125
96. Stute N, Trendelenburg U (1984) The outward transport of axoplasmic noradrenaline induced by a rise of the sodium concentration in the adrenergic nerve endings of the rat vas deferens. *Naunyn Schmiedeberg Arch Pharmacol* 327:124-132
97. Tani M, Neely JR (1989) Role of intracellular Na^+ in Ca^{++} overload and depressed recovery of ventricular function of reperfused ischemic rat hearts. *Circulat Res* 65:1045-1056
98. Tani M, Neely JR (1990) Na^+ accumulation increases Ca^{++} overload and impairs function in anoxic rat heart. *J Mol Cell Cardiol* 22:57-72
99. Tonnessen TI, Sandvig K, Olsnes S (1990) Role of Na^+/H^+ and $\text{Cl}^-/\text{HCO}_3^-$ antiports in the regulation of cytosolic pH near neutrality. *Am J Physiol* 258:C1117-C1126
100. Vandenberg JJ, Metcalfe JC, Grace AA (1993) Mechanisms of pH_i recovery after global ischemia in the perfused heart. *Circulat Res* 72:993-1003
101. Vaughan-Jones RD, Wu ML (1990) Extracellular H^+ inactivation of Na^+/H^+ exchange in the sheep cardiac Purkinje fibre. *J Physiol* 428:441-466
102. Wallert MA, Fröhlich O (1989) Na^+/H^+ exchange in isolated myocytes from adult rat heart. *Am J Physiol* 26:C207-C213
103. Wang HH, Katz RL (1965) Effects of changes in coronary blood pH on the heart. *Circulat Res* 17:114-122
104. Weiss ES, Hoffmann EJ, Phelps ME, Welch MJ, Henry PD, Ter-Pogossian MM, Sobel BE (1976) External detection and visualization of myocardial ischemia with ^{11}C -substrates in vitro and in vivo. *Circulat Res* 39:24-32
105. Weiss RG, Lakatta EG, Gerstenblith G (1990) Effects of amiloride on metabolism and contractility during reoxygenation in perfused rat heart. *Circulat Res* 66:1012-1022
106. Weissberg PL, Little PJ, Cragoe EJ, Bobik A (1989) The pH of spontaneously beating cultured rat heart cells is regulated by an ATP-calmodulin-dependent Na^+/H^+ antiport. *Circulat Res* 64:676-685
107. Williamson JR, Schaffer SW, Ford C, Safer B (1976) Contribution of tissue acidosis to ischemic injury in the perfused rat heart. *Circulation* 53:I3-I14
108. Winkler H, Apps DK, Fischer-Colbrie R (1986) The molecular function of adrenal chromaffin granules: established facts and unresolved topics. *Neuroscience* 18:261-290
109. Wollenberger A, Krause EG (1968) Metabolic control characteristics of the acutely ischemic myocardium. *Am J Cardiol* 22:349-359
110. Wright J, Maridonneau-Parini I, Cragoe EJ, Schwartz JH, Tauber AI (1988) The role of the Na^+/H^+ antiporter in human neutrophil NADPH-oxidase activation. *J Leukocyte Biol* 43:183-186
111. Zhao-Ping L, Chao-Shu T, Jing-Yi S (1991) Role of $\text{Na}^+/\text{Ca}^{++}$ exchange system in the pathogenesis of myocardial Ischemia-Reperfusion damage. *Science in China* 34:599-605

Received May 30, 1993
revision accepted August 8, 1993

Authors' address:

Wolfgang Scholz, Hoechst AG, SBU Cardiovascular Agents, H821, P.O. Box 800320, 65926 Frankfurt/M., FRG

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.